

# Commentary on Therapeutic Plasma Exchange to Treat COVID-19

Wen Lu\*

Department of Pathology, Brigham and Women's Hospital, Boston, United States

## Abstract

Severe acute respiratory syndrome coronavirus 2, the causative agent of coronavirus disease 2019, is capable of causing severe respiratory distress syndrome, cytokine release syndrome, multi-organ failure, and has been associated with many neurologic complications. No treatment has proven to be very effective. Therapeutic plasma exchange has been proposed and used. To date, randomized controlled trials have not been completed to show therapeutic benefit.

**Keywords:** Severe acute respiratory syndrome; Coronavirus; SARS-CoV-2

\*Corresponding author: Lu W

Department of Pathology, Brigham and Women's Hospital, Boston, United States.

 wlu5@bwh.harvard.edu

**Citation:** Lu W (2021) Commentary on Therapeutic Plasma Exchange to Treat COVID-19. Arch Clin Microbio Vol.12 S2: 148

**Received:** March 26, 2021; **Accepted:** April 09, 2021; **Published:** April 16, 2021

## Description

This March marked the one-year anniversary of the coronavirus disease 2019 (COVID-19) pandemic caused by Severe Acute Respiratory Syndrome Coronavirus 2 (SARS-CoV-2) [1]. Infection by SARS-CoV-2 may be asymptomatic or can cause severe disease affecting multiple organs including the lung, kidney, cardiovascular system, and gastrointestinal tract leading to Acute Respiratory Distress Syndrome (ARDS), Multi-Organ Failure (MOF), septic shock, and death [2]. In addition, SARS-CoV-2 has been associated with hypercoagulability, thromboembolic disease, Cytokine Release Syndrome (CRS), and a multitude of neurologic disorders.

Therapeutic Plasma Exchange (TPE) involves the removal of whole blood, its separation into components, with removal of plasma and return of non-plasma components along with replacement fluid (e.g., 5% albumin, Fresh Frozen Plasma [FFP]). There has been precedence for the use of TPE as adjunctive therapy in patients with viral infections who have failed conventional antiviral therapy alone [3]. For example, patients with ARDS due to avian influenza A (Hemagglutinin 7 Neuraminidases 9 [H7N9]) were treated with TPE and tandem continuous veno-venous hemofiltration. Patients with hemophagocytic lymphohistiocytosis and neurologic conditions thought to be triggered by H1N1 have also received TPE. Overall, outcome data after TPE in viral infections is sparse, variable, and inconclusive. The role of TPE in the treatment of neurologic complications associated with viral infection has not been established and lacks consensus guidelines. The only notable exceptions are in Guillain Barre Syndrome (GBS) and acute Myasthenia Gravis (MG) exacerbation. TPE is accepted as a first line therapy (category I indications) for both GBS and acute MG according to the 2019

American Society for Apheresis (ASFA) guidelines [4].

Otherwise, it is unclear that TPE is beneficial in the management of other neurological complications associated with SARS-CoV-2 infection and no high-quality data to show that TPE is helpful in COVID-19 related sepsis, MOF, and CRS. Nonetheless, TPE has been tried during the search for optimal management of patients with severe and complicated COVID-19. The theoretical rationale for TPE includes:

1. Attenuation of pro-inflammatory cytokines
2. Stabilization of the endothelial membrane
3. Normalization of hyperviscosity
4. Correction of aberrations in the coagulation pathway
5. Reduction of antifibrinolytic mediators
6. Fibrin degradation products
7. SARS-CoV-2 virus

These mechanisms, however, remain to be proven. In part because given the novel nature of SARS-CoV-2, much of the pathogenic pathways that lead to the complications of COVID-19 are not well understood. A final consideration is that TPE removes plasma constituents non-selectively. Therefore, removal of protective host defense proteins, anti-inflammatory mediators, anti-SARS-CoV-2 neutralizing antibodies, and potentially effective drugs (e.g, antivirals, anti-inflammatory) may actually hamper resolution of the infection [5-12].

The case reports and case series published on the use of TPE as adjunctive therapy for COVID-19 are inherently limited in their study design [12-15]. No prospective randomized controlled trials have been completed to establish that TPE is effective

therapy for COVID-19. While generally safe and well tolerated, as with any medical intervention, TPE is not without risk. Routine complications of TPE associated with central line placement, line infection, and hypocalcemia apply for COVID-19 patients. While complications of severe hypocalcemia (hypotension, arrhythmia, and QT prolongation) are rare, they are of particular concern in critically ill patients with preexisting septic shock and MOF. Replacement without plasma can cause hypokalemia and decreased coagulation factors. Replacement with plasma puts patients at risk of adverse events associated with transfusion. An additional consideration is the risk of clotting during the procedure which may prematurely interrupt the procedure and lead to acute blood loss in very sick patients.

If TPE is to be performed in patients with COVID-19, it should be done as part of one of the registered clinical trials. Completion of these trials is the only way to determine the safety and efficacy of TPE in COVID-19.

## References

1. Coronavirus Disease 2019 (COVID-19) Situation Report – 199. World Health Organization. Accessed March 23, 2021.
2. Cevik M, Bamford CGG, Ho A (2020) COVID-19 pandemic-a focused review for clinicians. *Clin Microbiol Infect* 26: 842-847.
3. Lu W, Kelley W, Fang DC, Joshi S, Kim Y, et al. (2021) The use of therapeutic plasma exchange as adjunctive therapy in the treatment of coronavirus disease 2019: A critical appraisal of the current evidence. *J Clin Apheresis*.
4. Padmanabhan A, Smith LC, Aqui N (2019) Guidelines on the use of therapeutic apheresis in clinical practice – Evidence-based approach from the writing committee of the American Society for Apheresis: The eighth special issue. *J Clin Apheresis* 34:171-354.
5. Swol J, Lorusso R (2020) Additive treatment considerations in COVID-19-The clinician's perspective on extracorporeal adjunctive purification techniques. *Artif Organs* 44: 918-925.
6. Keith P, Day M, Perkins L, Moyer L, Hewitt K, et al. (2020) A novel treatment approach to the novel coronavirus: An argument for the use of therapeutic plasma exchange for fulminant COVID-19. *Crit Care* 24:128.
7. Zhang L, Zhai H, Ma S, Chen J, Gao Y (2020) Efficacy of therapeutic plasma exchange in severe COVID-19 patients. *Br J Haematol* 190: 181-183.
8. Maier CL, Truong AD, Auld SC, Polly DM, Tanksley CL, et al. (2020) COVID-19-associated hyperviscosity: a link between inflammation and thrombophilia? *Lancet* 395: 1758-1759.
9. Ahmed S, Zimba O, Gasparyan AY (2020) Thrombosis in Coronavirus disease 2019 (COVID-19) through the prism of Virchow's triad. *Clin Rheumatol* 39:2529-2543.
10. Turgutkaya A, Yavasoglu I, Bolaman Z. The Application of plasmapheresis For Covid-19 patients. *Ther Apher Dial* 25: 248-249.
11. Alexander D, Truong AD, Sara C Auld SC, Nicholas A, et al. (2020) Therapeutic plasma exchange for COVID-19-associated hyperviscosity. *Transfusion*.
12. Gucyetmez B, Atalan HK, Sertdemir I, Cakir U, Telci L (2020) COVID-19 study group. therapeutic plasma exchange in patients with COVID-19 pneumonia in intensive care unit: A retrospective study. *Crit Care* 24:492.
13. Khamis F, Al-Zakwani I, Al Hashmi S (2020) Therapeutic plasma exchange in adults with severe COVID-19 infection. *Int J Infect Dis* 8.
14. Adeli SH, Asghari A, Tabarraii R (2020) Therapeutic plasma exchange as a rescue therapy in patients with coronavirus disease 2019: A case series. *Pol Arch Intern Med* 130: 455-458.
15. Faqihi F, Alharthy A, Alodat M, Kutsogiannis DJ, Brindley PG, et al. (2020) Therapeutic plasma exchange in adult critically ill patients with life-threatening SARS-CoV-2 disease: A pilot study [published online ahead of print, 2020 Jul 31]. *J Crit Care* 60: 328-333.