

DOI: 10.36648/1989-8436.11.3.113

The Biological Activity of Aqueous and Methanolic Extracts of *Juglans regia* on Yeasts and Pathogenic Bacteria

Furdos N Jafer and Lina A Naser*

Biology Department, Science College, Basrah University, Iraq

*Corresponding author: Lina A Naser, Biology Department, Science College, Basrah University, Iraq, Tel: 07705698101; E-mail: lina1977_abbas@yahoo.com

Received date: October 21, 2019; Accepted date: June 05, 2020; Published date: June 12, 2020

Citation: Jafer FN, Naser LA (2020) The Biological Activity of Aqueous and Methanolic Extracts of *Juglans regia* on Yeasts and Pathogenic Bacteria. Arch Clin Microbiol Vol. 11 No. 3:113

Copyright: © 2020 Jafer FN, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Abstract

Background: The plant extract can be used as a natural alternative to control of pathogenic yeasts and bacteria in general and to reduce the risk of antibiotic-resistant pathogens.

Methods: Diffusion agar method was used to examine the antifungal and antibacterial activity of both aqueous and methanolic *Juglans regia* extracts. For yeasts, The MIC values were determined by the standard serial dilution assay using serial dilutions of the aqueous and methanolic extracts (300,200,100, 50, 25) µg/L). The MIC for bacteria was done using Muller-Hinton agar. The concentrations of the extracts tested were (500, 250, 100, 50) µg/ml.

Results: *Juglans regia* extracts showed activities against *Malassezia furfur* yeast, Methanolic extract exhibits more activity than aqueous extract. The Minimum Inhibition Concentrated rate (MIC) for *Malassezia furfur* methanolic extract was 75 µg/ml. The extracts also examined on *Candida albicans* opportunistic yeast in the mouth and found to be more active than *M. furfur* were MIC 50 µg/ml. The biological activity of aqueous and methanolic extracts were also examined against six pathogenic bacteria isolated from patients suffering from different diseases from several hospitals in Basrah, including *Staphylococcus aureus*, *Escherichia coli*, *Salmonella typhi*, *Proteus mirabilis*, *pseudomonas aeruginosa* and *Vibrio parahaemolyticus*. The Methanolic extract showed an effective effect in inhibiting all bacterial species compared with the aqueous extract.

Conclusion: This study provides the possibility of using a natural plant such as *Juglans regia* as antimicrobial agent for many pathogenic yeasts and bacteria.

Introduction

Some fungi found on human skin and scalp, such as *Malassezia*, and are found in some human cavities such in the mouth and vagina such as *Candida*. *Malassezia* is yeast that is naturally found on the skin of healthy human and warm-blooded animals and can become a pathogen and cause skin diseases such as Pityriasis versicolor [1-4].

This yeast tends to fat, therefore called lipophilic and all sex-related species of the seven species considered to be lipophilic, except *M. pachyderms*. Therefore, the medium used to isolate should contain fatty acids [5-7]. Olive oil is one of the most common fats used to isolate this yeast. The most important characteristic of this yeast is the possession of the enzyme Lipase [8]. *Candida* is bivalent yeast that is co-existent in warm-blooded animals, including humans. It colonizes the mucous surfaces of the mouth, vagina, and digestive tract. It can cause various infections, according to the host. The infection caused by *Candida* is called Candidiasis and was two types oral (Thrash) and vaginal candidiasis. *Candida* is dimorphic yeast it has two phases, yeast phase, and the filament phase and is transformed from one phase to another depending on PH and temperature and the presence of other compounds such as Acetylglucosamine [9-12].

Some primitive methods of decoration are used, such as the use of *Juglans regia* to whiten the teeth and apply dye to the lips [13].

The walnut trees live in northern Iran, the Caucasus, Baluchistan, and Armenia, and these perennial trees date back to 500 years [14]. It was used as an antifungal agent especially in the treatment of ringworm infections for the first time in 1950 by the world Dorian. The extract successfully used in inhibiting yeast growth *Candida albicans* [15].

The plant extract can be used as a natural alternative to control of pathogenic bacteria in general and to reduce the risk of antibiotic-resistant pathogens in particular [16].

In a study carried out by Sharafati [17], the *Juglans regia* extract showed a wide spectrum effect against bacteria such

Keywords: *Juglans regia*; Activity; Antifungal; Antibacterial

as *Staphylococcus aureus*, *Streptococcus mutans*, *Escherichia coli*, *Pseudomonas aeruginosa*.

The discovery of antibiotics in the last century led to a decrease in disease rates and deaths due to infectious diseases, but the inappropriate use of these antibiotics resulted from the emergence of groups of resistant bacteria that threaten human health, especially with immunocompromised patients, creating the need to research new types of natural anti-microbial substances to control on different diseases and has led to an increase in interest in medicinal plants where 25-50% of the current pharmaceuticals derived from plants because chemically manufactured drugs have side effects, while there is no such effects in medical alternatives such as plant extracts [18].

Therefore, the aim of this study was to study the antimicrobial effect of *Juglans regia* extract (aqueous and Methanolic) on yeasts and pathogenic bacteria isolated from patients suffering from different diseases in Basrah hospitals.

Materials and Methods

Preparation of aqueous and methanolic *Juglans regia* extracts: 20 gm of the powdered dry bark of *Juglans regia* were weight separately in different flasks, one of them percolated with 200 ml of methanol this flask was proper sealed with aluminum foil and left with stirring magnetic stirrer for 24 hours at room temperature. The solution was then filtered using a funnel filled in a filter paper and the methanol extracts collected in Petri dishes and dried at room temperature. The above procedure was repeated with the use of 200 ml of distilled water to prepare the aqueous extract [19].

Isolation of yeasts

Malassezia yeast isolated from person with pityriasis versicolor by used of skin scraping that cultured in Sabouraud Dextrose Agar (SDA) medium supplemented with 20 ml of olive oil per liter, 0.01 cyclohexamide and 10 drops of tween 20. The isolated *Malassezia* classified as *Malassezia furfur* While *Candida* yeast isolated from person with mouth candidiasis infection.

Oral swab used and cultured in (SDA) both yeasts incubated in 32°C the isolated candida classified as *Candida albicans* by using of germ tube test.

Study the effect of alternative oils on the *Malassezia* yeast growth

Appropriate medium for the growth of *Malassezia* is the added to some oils communally olive oil and Tween 80 as mentioned in the introduction, but in this experiment has been replaced olive oil with some oils such as sunflower oil and castor oil and olive oil treatment compared to add to use a combination of the three types as an independent treatment, the previous oils were added by 20 ml per liter of SDA media. Each of the cultivars was poured into the dishes. The yeast was

then planted on the medium in a streaking manner, taking into consideration three replicates per treatment and incubated at 32°C.

Antifungal activity test

Diffusion agar method was used to examine the antifungal activity of both aqueous and methanolic extracts toward *Malassezia furfur* and *Candida albicans* using (SDA) medium, 10 µl of each yeast suspension add on the surface of the medium then spread with sterile L-shape glass rod, after drying, two 5 mm diameter pores was made in the center of plates, 100 µl of both crud extracts was add separately in the pores of plates with used of control treatment for each yeast, cultures were incubated at 32°C for 2 days after incubation period the diameter of the inhibition zones were evaluated in millimeters [20].

The minimum inhibitory concentration (MIC) test for yeast

The MIC values were determined by the standard serial dilution assay using serial dilutions of the aqueous and methanolic extracts (300,200,100, 50, 25), µg/L. The suspensions of yeasts were prepared and standardized to a turbidity equivalent to that of 0.5 MacFarland scale (1×10⁶ cfu/ml). 100 µl of yeast suspension add on the surface of the media then media incubated in 32°C, The MIC values in this assay were indicated by the absence of yeast growth at the minimal concentration of the extract.

Antibacterial activity test

The Muller Hinton agar was used for the purpose of measuring the biological efficacy of six types of pathogenic bacteria *Staphylococcus aureus*, *Escherichia coli*, *Salmonella typhi*, *Proteus mirabilis*, *Pseudomonas aeruginosa* and *Vibrio parahaemolyticus* were isolated from patients suffering from different diseases admitted to Al Fayhaa General Hospital and Al Shifa General Hospital in Basrah. The well-diffusion methods were used by Bansode and Chavan [21] 0.1 ml of the bacteria was distributed in the Muller Hinton agar dish using a sterile swab and then holes were made in the center of the plates using a cork borer. 100 µl of both aqueous and methanolic crude extracts were placed separately in each hole. One distilled water was placed in a single hole as a control. The dishes were then left for half an hour after being placed in the incubator for 24 hours at 37°C. The effectiveness of the two extracts was determined by measuring the diameter of the inhibition zone formed around the hole mm.

The Minimum Inhibitory Concentration (MIC) test for bacteria

The MIC of different concentrations of Methanolic and aqueous *Juglans regia* extracts against tested bacterial isolates were done using Muller-Hinton agar. The concentrations of the extracts tested were (500, 250, 100, 50) µg/ml, the assays were performed three times for each bacterium. Plates were

inoculated with 0.1 ml of the bacterial suspension was distributed in the Muller Hinton agar dish using a sterile swab and then holes were made using a cork borer. Different extract was placed in each hole. The plates then left for half an hour after being placed in the incubator for 24 hours at 37°C. The highest dilution of extract that showed no visible bacterial growth per hole was considered as MIC Standards.

Results

Preparation of aqueous and methanolic Juglans regia extracts

After the preparation of the extract it was observed that the aqueous extract is characterized by viscous nature while the methanolic extract appeared in the crystalline nature due to the difference in the type of chemical compounds extracted from each one.

Isolation and Diagnosis of Yeast

After the *Malassezia Yeast* isolate from the Skin infection pityriasis versicolor the colony was raised and smooth initially and get dry and wrinkle in time the color of colony was creamy to a slightly yellowish-colored. The microscopic examination indicated that it represented the *M. Furfur*.

Species *M. furfur* is an ellipsoids yeast with small collarets and contains a bud or scar from one side as a characteristic feature.

The other yeast isolate was *Candida albicans*, which had a cream glistening and sometime waxy surface is smooth. The microscopic examination of *Candida albicans* was small oval with single budding sometime this yeast was give positive germ tube test.

Study the effect of alternative oils on the Malassezia yeast growth

Experiment showed that the best growth of *Malassezia furfur* was in the medium containing olive oil and sunflower oil. Either the medium with other treatment of castor oil appeared weak growing of *Malassezia furfur*, mixing oils as results of castor oil (Figure 1).

Antifungal activity test

Both aqueous and methanolic extracts showed effectiveness in inhibiting the growth of both yeasts. However, the effect of the extracts was higher with respect to the yeast *C. albicans* compared with the yeast *M. furfur*. The diameters of the inhibitory zone were 20 mm and 25 mm respectively for the aqueous and methanolic extracts. *C. albicans* (45) mm for aqueous extract and (3) mm for methanolic extracts (Table 1 and Figure 2).




Figure 1 The growth of *Malassezia furfur*, on the medium containing different oils A-Olive oil B-Sunflower oil C-Castor oil D-Oil mixture

Table 1 The diameter of the inhibition zones of extracts toward yeasts

Inhibition zone diameter(mm)		
Yeast	Aqueous extract	Methanolic extract
<i>M. furfur</i>	20	25
<i>C. albicans</i>	45	30



Figure 2 Indicate the diameters of the inhibition zone of *C. albicans* in the effect of the extracts A-Control B-Aqueous extract C-Methanolic extract

The MIC test for yeast

The concentrations of 200 and 300 µg/ml showed a lack of growth for both extracts, MIC for *C. albicans* yeast 50 µg/ml in the aqueous extract and 75 µg/ml in the Methanolic extract.

MIC for *M. furfur* yeast in the treatment of the aqueous extract was 100 µg/ml and 75 µg/ml in methanolic extract obtained (Table 2).

Table 2 MIC of extracts toward yeasts.

MIC value ($\mu\text{g/ml}$)		
yeast	Aqueous extracts	Methanolic extracts
<i>M. furfur</i>	100	75
<i>C. albicans</i>	50	75

Antibacterial activity test

Two positive and negative Gram bacteria were selected to determine the effect of *Juglans regia* extracts on them.

Methanolic extract showed a clear and almost fatal effect on most bacterial species (**Table 3**). *Staphylococcus aureus* showed the most sensitive germs, followed by *Salmonella typhi*.

Table 3 Results of the inhibitory effect and diameters of the inhibition zones in (mm) for the aqueous and Methanolic extract against 6 bacterial isolates.

Pathogenic Bacteria	Diameters of mm inhibition zone	
	Alcoholic extract Mm/mg	Water extract Mm/mg
<i>Staphylococcus aureus</i>	22.5	15.5
<i>Pseudomonas aeruginosa</i>	7	8
<i>Vibrio parahaemolyticus</i>	12.5	R
<i>Salmonella typhi</i>	20	10
<i>Proteus mirabilis</i>	11	10
<i>Escherichia coli</i>	17.5	15.5

Where R means resistance

The MIC for pathogenic bacteria

Concentrations showed 50 micrograms/ml growth loss for all bacterial pathogens in both extracts. MIC for *Vibrio parahaemolyticus* and *Pseudomonas aeruginosa* 500 $\mu\text{g/ml}$ in the water extract were 250 $\mu\text{g/ml}$ in the alcohol extract.

MIC for *Staphylococcus aureus* showed a concentration of 50 $\mu\text{g/ml}$ growth in the treatment of aqueous and methanolic extract and **Table 4** shows this.

Table 4 The MIC of the pathogenic bacteria under study towards aqueous and alcohol extract.

Bacteria	500		250		100		50	
	C	W	C	W	C	W	C	W
<i>Staphylococcus aureus</i>	+	+	+	+	+	+	-	-
<i>Pseudomonas aeruginosa</i>	+	-	-	-	-	-	-	-
<i>Escherichia coli</i>	+	+	+	+	+	-	-	-
<i>Vibrio parahaemolyticus</i>	+	-	-	-	-	-	-	-
<i>Salmonella typhi</i>	+	+	+	-	-	-	-	-
<i>Proteus mirabilis</i>	+	+	+	-	-	-	-	-

Discussion

The experiment demonstrated that olive oil and sunflower oil were comparable in their effectiveness to support the growth of *M. furfur* compared to castor oil [8] because it is high in density and complex structure which is difficult to digest into simple fatty acids by Lipase enzyme and contains a protein called Resin inhibits protein building in the cell.

The extract of *Juglans regia* was tested against *M. furfur* as tested for yeast *C. albicans*, and extract is used as a cosmetic to beautify the lips and whiten teeth in the mouth [13].

The use of plant extracts is still common due to the resistance of many types of yeast to antifungal agents. The extract of *Artemisia abrotanum* has been successfully used to inhibit the growth of yeast *M. furfur* and *C. albicans* [22].

Essential oils derived from orange and lemon has been found to be effective against the fungus *M. Furfur*, which causes varicose veins [23].

Juglans regia is used as an extract in the inhibition of the growth of many skin fungi and with high efficiency such as *Epidermophyton floccosum* and *Trichophyton mentagrophytes* and *C. albicans*. Al-Khawaja [13] noted that the use of *Juglans regia* plant in the mouth to clean the teeth and adornment increases the acidic function of the PH saliva, which leads to an antimicrobial action. The experiment proved superior to the aqueous than methanolic extract in the antimicrobial activity experiment on *C. albicans* and has a higher MIC, this is due to the ability of water to pull many compounds and active ingredients (alkaloids, tannins and amino acids) from their plant source [24]. While the result was reflected in the yeast of *M. furfur* in the effectiveness of antifungal activity and MIC, the methanolic extract was more effective in inhibition of the growth this is because to the fact that mehanol is able to extract flavonoids that may affect this yeast [25]. This is agreement with Qadan [26] found that phenolic compounds such as flavonoids have anti-fungal effects. Buttery noted that *Juglans regia* is containing terpenes and hydrocarbons Esters that have antifungal and antioxidants action such as phenolic compounds.

It is recognized that antibacterial resistance to antibiotics has been rising to dangerous levels around the world. Antibiotic resistance leads to prolonged hospital stay, high medical costs and increased mortality. New resistance mechanisms are emerging and spreading globally and threaten our ability to treat common infectious diseases [27-33].

Conclusion

Antibiotic resistance increases and spreads in cases where the over-the-counter purchase of antibiotics is considered for human or animal use. All these things called for rethinking the return to natural resources.

The results of the test of the efficacy of aqueous and methanolic extracts of *Juglans regia* as an anti-bacterial agent showed that gram-negative bacteria and gram-positive bacteria were affected by the both extracts used.

Juglans regia methanolic extract showed zone of inhibition against all tested bacterial strains, while the aqueous extract was active with relatively smaller inhibition zones, which is consistent. *S. aureus* and *Salmonella typhi* showed that they were the most sensitive species for the effect of methanolic extracts for the extract of water of the *Juglans regia*, bacteria *S. aureus* showed high sensitivity while *Salmonella* appeared less sensitive.

The results showed that bacteria *Vibrio parahaemolyticus* were sensitive to the alcohol extract, while the aqueous dermis showed no efficacy against them.

Financial Support and Sponsorship

Nil.

Conflicts of Interest

There are no conflicts of interest.

References

1. Ahmad S, Wahid MA (1973) Fungistatic action of *Juglans* l3: 436-438.
2. AL-Khawajah AM (1997) Studies on the antimicrobial activity of *Juglans regia*. The Amer of chires. Medic (AJCM) 25: 170-180.
3. Alkhawajah A (2012) Studies on the Antimicrobial Activity of *Juglans regia*. The American Journal of Chinese Medicine 25.
4. Bansode DS, Chavan MD (2013) Evaluation of antimicrobial activity and phytochemical analysis of papaya and pineapple fruit juices against selected enteric pathogens. International Journal of Pharma and Bio Sciences 4: B1176-B1184.
5. Brodin K, Alahyar H, Heduer T, Faergemanr J (2007) Invitro Activity of Artemiisia abrotanum extract aginst malassezei spp. *Candida albicans* and staph. aureus. aceta Dermat. venergeologica, 87: 540-542.
6. Buttery RG, Fiat R, Mon TR, Anclling CL (1986) Identification germacrene Din walnut and fig leat volatiles. TA food Chem 34: 820-822 .
7. Chopra RN, Copra IC, Handa KL, Kapur LD (1958) Indigenons drugsof India, 2nd ed. VN Dhur tsors, private Ltd. Calcutta.
8. Cutler JE (1991) Putative virulence factor of candida albicans. Anhu revmicro boil 45: 187-218.
9. Dismukes W, Pappas P, Sobel J (2003) Clinical Mycology Oxford University Press, 504.
10. Dorlons WA (1951) Quotel in Americn illust voted medical dictionary, 22nd ed. Saunders and Co. Philadelphia.
11. Grim show J (1976) Despsides, hydrolysable tannins, lignans, lignin and Humic acid. coffeye 11 Pewt D. elsever scientific puplishemy co. Amsterdam, Netherlands.
12. Gueho Midgley G, Guollot J (1996) The genus *Malassezia* with description of four new species Antoine van Leeuwenhoek 96: 337-557.
13. Gueho E, Boekhout T, Ashbee guillot J, vanbelkun A, faergemann J (1998) The role of *Malassezia* species in the ecology of human skin as pathogen. Med Mycol 36: 220-229.
14. Gupta PG, Birdi TJ (2017) Development of botanicals to combat antibiotic resistance. J Ayurveda Integr Med 8: 266-275.
15. Handa SS, Khanja SPS, Longo G, Rakesh DD (2008) Extraction Technologies for medicinal and Aromatic plants, International Centre for Science and High Technology, Trieste 21-25.
16. Harborn JB (1984) Phytochemical Methods chpman and Hall, London, UK.
17. Ingham E, Cunningham A (1993) *Malasszia furfur*. J Med and Vet Mycol. 31: 265-288.
18. Leberer E, Zigelbauer K, Schmidt A (1997) Virulance and hyphat formation of candida albicans vequire the stezop-like protion kinase caclou p. current biology 7: 539-546.
19. Lo HJ, Kohler JDI, Domenico BL, Domenico B, Loebenberge (1990) Non filamentous *C. albicons* mutants are avivule cell 90: 939-949.
20. Lucia KH, Cecilia MA, Pedro HF (2003) Antimicrodial Activity of huptis ollaliforia towaral dermato phytes. Men Inst of Fwado Cruze Riode Janerio 98: 963-965.

21. Midgley G, Gueho E, Guillot J (1997) Diseases caused by *Malassezia* spp. in: Medical Mycology. (eds.) Hay RJ and Ajello L Arnold Publ, London 4: 201-211.
22. Qadan FA, Thewaini DA, Ali RA, Matalka KZ (2005) The Antimicrobial activity of *Psidium Gnajawa* and *Juglans regia* leaf extract to ocnedeveloping organisms amrec. Jour Chin Med (AJCM) 33: 197-205.
23. Ran Y, Yoshiike T, Ogawa H (1993) Lipase of *Malassezia* same properties and their relationship to cell growth. J Med veta myccel 31: 77-85.
24. Rossiter SE, Fletcher MH, Wuest WM (2012) Natural Products as Platforms to Overcome Antibiotic Resistance. Chem Rev 117: 12415-12474.
25. Ryley JF, Ryley MG (1995) *Candida albicans* - do mycelia matter? Jmed Vetmycol 28: 225-239.
26. Salamat F, Keivani S, Amin G (2006) Evolution of *Juglans regia*. pericrp on antifng susceptibility with both dilution method. Med Scen Jour of Islamic Azad vniv. 16: 201-205.
27. Sharafati Chaleshtori R, Sharafati-Chaleshtori F, Rafeian-Kopaei M (2011) "Biological characterization of Iranian walnut (*Juglans regia*) leaves," Turkish Journal of Biology 35: 635-639.
28. Sharma R, Sharma G, Sharma M (2012) Anti-*Malassezia* furfur activity of essential oil against causal agent of pityrias versicolor disease African Jaur. Of Pharmacy and pharmacology vol 6: 979-983.
29. Spooner DF, Sykes G (1972) Laboratory assessment of antibacterial activity. In: nmethodsin microbiology, Norris JR, Ribbors DW (eds) 7: 211-278.
30. Sytykiewicz H, Chrzanowski G, Czerniewic P, Leszczyns B (2015) Antifungal activity of *Juglans regia* (L.) leaf extracts against *Candida albicans* isolates. Pol J Environ Stud 24: 1339-1348.
31. Tiwari R, Tiwari G (2011) Use of antibiotics: from preceding to contemporary. Scholars Res J 1: 59-68.
32. Yarrow D, Ahearn D (1984) *Malassezia* Bailion in the yeasts. A taxonomic study. (3rdedn) kreger-van Rij. NJW Amsterdam Elsevier science publishers Bv 882-885.
33. Zakavi F, Hagh L, Daraeighadikolaei A, Sheikh A, Daraeighadikolaei A, et al. (2013) Antibacterial Effect of *Juglans Regia* Bark against Oral Pathologic Bacteria. International Journal of Dentistry P: 5.